



PrimeWay Soil DNA Extraction

Wash Buffer 52 Wash Buffer 51

PrimeWay Soil DNA Extraction Kit is a reliable kit that is used to isolate genomic DNA from various type of soil sample, manure & water sample.


Soil

- ➤ General soil
- ➤ Low microbial diversity soil

Manure **Water Sample DNA Clean-up**


Both mechanical and chemical lysis methods are used for maximum extraction efficiency and DNA yield. This kit can efficiently remove abundance of humic substances and pigments which affect downstream processes such as PCR. Besides soil sample, it is also suitable for other sample types including animal manure, worm compost and water. The purified DNA is suitable for PCR, southern blot, enzyme digestion, amplicon sequencing, etc.

Performance Review Soil




DNA isolated from 500 mg various types of soil.

PCR Amplification



50 ng of DNA are analysed with 1% agarose gel.



Successful PCR amplification indicates the extracted DNA is free from PCR inhibitors. 1 µL PCR product is analysed with 1% agarose gel.

OG: Organic soil OP: Oil Palm estate soil FS: Forest soil VG: Vegetable bed soil

PD: Paddy soil


More Details






Comparison Data

DNA Yield, Purity & Integrity for General Soil



DNA isolated from 200 mg soils.




50 ng of DNA are analysed with 1% agarose gel.

Soil Type OG: Organic soil FS: Forest soil PD: Paddy soil <u>Brand</u>

PW: PrimeWay
M: Brand M
PL: Brand PL
Q: Brand Q

PCR Amplification



Successful PCR amplification (ITS2) indicates the extracted DNA is free from PCR inhibitors. 1 μ L PCR product is analysed with 1% agarose gel.












DNA Yield, Purity & Integrity for Low Soil Microbial Diversity




DNA isolated from 200 mg clay. PrimeWay kit able to extract DNA from soil with low microbiome diversity.

1kb PL Q

 $50\ ng$ of DNA are analysed with 1% agarose gel.

PCR Amplification



Successful PCR amplification (ITS2) indicates the extracted DNA is free from PCR inhibitors. 1 µL PCR product is analysed with 1% agarose gel.


<u>Brand</u>

PW: PrimeWay M: Brand M PL: Brand PL Q: Brand Q


Manure

Performance Review

DNA Yield, Purity & Integrity




DNA isolated from 500 mg of cow manure & worm compost.



50 ng of DNA are analysed with 1% agarose gel.

CM: Cow Manure WC: Worm Compost

PCR Amplification



Successful PCR amplification (ITS2) indicates the extracted DNA is free from PCR inhibitors. 1 µL PCR product is analysed with 1% agarose gel.









Comparison Data

DNA Yield, Purity & Integrity for Manure




DNA isolated from 100 mg of cow manure & worm compost.



50 ng of DNA extracted are analysed with 1% agarose gel.

Soil Type CM: Cow Manure WC: Worm Compost <u>Brand</u> PW: PrimeWay M: Brand M PL: Brand PL Q: Brand Q

PCR Amplification



Successful PCR amplification (ITS2) indicates the extracted DNA is free from PCR inhibitors. 1 µL PCR product is analysed with 1% agarose gel.









Performance Review Water

DNA Yield, Purity & Integrity




DNA isolated from various type of water sample with 0.2 μm membrane filter.



50 ng of DNA from water samples are analysed with 1% agarose gel.

Water Sample Type LK: Lake water WS: Waste water SW: Sea water

PCR Amplification



Successful PCR amplification (ITS2) indicates extracted DNA is free from PCR inhibitors. 1 μL PCR product is analysed with 1% agarose gel.








DNA Clean-up


Performance Review

DNA Recovery & Purity after DNA Clean-up



DNA clean-up of 30 µL DNA containing pigments.

Absorbance 320, Humic Acid Detection



Before



Light yellowish pigment is observed in the DNA

After



The pigments in the DNA are removed

